

COMPARING CHEBYSHEV POLYNOMIALS AND ADOMIAN DECOMPOSITION
METHOD IN SOLVING NONLINEAR VOLTERRA INTEGRAL EQUATIONS
OF SECOND KIND

SITI AMINAH BINTI MOHAMAD SAPAWI

UNIVERSITI TEKNOLOGI MALAYSIA

COMPARING CHEBYSHEV POLYNOMIALS AND ADOMIAN
DECOMPOSITION METHOD IN SOLVING NONLINEAR VOLTERRA
INTEGRAL EQUATIONS OF SECOND KIND

SITI AMINAH BINTI MOHAMAD SAPAWI

A dissertation submitted in partial fulfillment of the
requirements for the award of the degree of
Master of Science (Mathematics)

Faculty of Science
Universiti Teknologi Malaysia

JUNE 2014

Specially dedicated to my beloved parents,
Mohamad Sapawi bin Ramli and Zanariah bt Mahmud

ACKNOWLEDGEMENT

Alhamdulillah, thanks to Allah because by His will, I manage to complete this dissertation. This dissertation means a lot to me, since I learn more new knowledge. I wish to express my sincere appreciation to my supervisor, Assoc. Prof. Dr Munira Ismail for encouragement, guidance, and critics. Without her continued support and instruct, this report would not been completed.

Besides, I would like to take this opportunity to thanks my family who always support me and my friends that willing to teach and assists me to complete this dissertation. They really supported me and played an important role in the completion of this dissertation.

ABSTRACT

The nonlinear integral equations are usually difficult to solve analytically and in many cases, it is required to obtain the approximate solutions. The nonlinear Volterra integral equation of second kind is one of them. This dissertation compares two methods that are used in order to solve nonlinear Volterra integral equation of second kind. Those are Chebyshev polynomials and Adomian decomposition method. The Chebyshev polynomials are developed to approximate the solution of linear and nonlinear Volterra integral equations. While, Adomian decomposition method, is a method that can be applied directly for all type of linear and nonlinear integral equations and maintain high accuracy of numerical solution. Hence, the best method is picked based on the absolute error that will be compared with the exact solution.

ABSTRAK

Persamaan kamiran tidak linear kebiasaannya sukar diselesaikan secara analitik dan untuk menyelesaikannya memerlukan penyelesaian anggaran. Persamaan kamiran Volterra tidak linear merupakan salah satu darinya. Penyelidikan ini membandingkan dua kaedah untuk menyelesaikan Persamaan kamiran Volterra tidak linear iaitu kaedah polynomial Chebyshev dan kaedah penguraian Adomian. Kaedah polynomial Chebyshev dibentuk untuk menganggarkan penyelesaian persamaan kamiran Volterra linear dan tidak linear. Manakala, kaedah penguraian Adomian merupakan kaedah yang boleh digunakan secara langsung samaada pada persamaan kamiran tidak linear atau linear dengan mengekalkan ketepatan daripada penyelesaian berangka. Kaedah yang paling baik dipilih dari penyelidikan ini berdasarkan ralat mutlak apabila dibandingkan dengan penyelesaian tepat.